

Appendix C: List of included studies

1. Ahmed SS, Onyambu CK, Omamo E, Odhiambo A. Appropriateness of imaging modality choice by doctors at the Kenyatta National Hospital's Accident and Emergency Department. *S Afr J Radiol.* 2022;26(1):a2367. Available from: <https://doi.org/10.4102/sajr.v26i1.2367>
2. AlJaroudi W, Isma'eel H, El Merhi F, Assad T, Hourani M. (2015). Appropriateness and diagnostic yield of cardiac magnetic resonance imaging from a tertiary referral center in the Middle East. *Cardiovasc Diagn Ther.* 2015;5(2):88-97. Available from: <https://doi.org/10.3978/j.issn.2223-3652.2014.11.01>
3. Alimohammadi H, Zareh Shahamati S, Karkhaneh Yousefi A, Safarpour Lima B. (2021). Potentially inappropriate brain CT-scan requesting in the emergency department: A retrospective study in patients with neurologic complaints. *Acta Bio Medica : Atenei Parmensis*, [online] 92(5), p.e2021302. Available from: <https://doi.org/10.23750/abm.v92i5.10209>
4. Arjona M, Toldo JMP, Queiroz NCM, Pedroso JL, de Carvalho Campos Neto G, Barsottini OGP, et al. A real-world study of cerebral 99mTc-trodat-1 single-photon emission computed tomography (SPECT) imaging of the dopamine transporter in patients with Parkinson disease from a tertiary hospital in Brazil. *Med Sci Monit.* 2020;26:e925130. Available from: <https://doi.org/10.12659/msm.925130>
5. Baiguissova D, Laghi A, Rakhimbekova A, Fakhradiyev I, Mukhamejanova A, Battalova G, et al. An economic impact of incorrect referrals for MRI and CT scans: a retrospective analysis. *Health Sci Rep.* 2023;6(3):e1102. Available from: <https://doi.org/10.1002/hsr2.1102>
6. Becker J, Jenkins LS, De Swardt M, Sayed R, Viljoen M. Appropriateness of computed tomography and magnetic resonance imaging scans in the Eden and Central Karoo districts of the Western Cape Province, South Africa. *S Afr Med J.* 2014;104(11):762. Available from: <https://doi.org/10.7196/samj.8158>
7. Demeke E, Mekonnen A. Appropriateness of head CT scans at Tikur Anbessa Specialized Hospital, Ethiopia. *Ethiop J Health Sci.* 2022;32(2):359-68. Available from: <https://doi.org/10.4314/ejhs.v32i2.17>
8. Denjagić A. Justification of radiological procedures algorithm adjustment in diagnosis of lower back pain cause at University Clinical Center Tuzla. *Acta Medica Saliniana.* 2019;49(1):33-40. Available from: <https://doi.org/10.5457/ams.v49i1.489>
9. Dos Santos MA, Santos MS, Tura BR, Félix R, Brito ASX, De Lorenzo A. Budget impact of applying appropriateness criteria for myocardial perfusion scintigraphy: the perspective of a developing country. *J Nuc Cardiol.* 2016;23(5):1160-5. Available from: <https://doi.org/10.1007/s12350-016-0505-4>
10. Edzie EKM, Dzefi-Tetty K, Gorleku PN, Brakohiapa EK, Botwe BO, Amankwa AT, Idun EA, Kusodzi H, Asemah AR. Audit of the appropriateness of the indication for obstetric sonography in a tertiary facility in Ghana. *Pan Afr Med J.* 2021;40(35). Available from: <https://doi.org/10.11604/pamj.2021.40.35.26349>
11. Fouche PE, Jenkins L, Vermeulen A. Appropriateness of computed tomography and magnetic resonance imaging scans in a rural regional hospital in South Africa: a 6-year follow-up study. *S Afr Med J.* 2020;111(1):46. Available from: <https://doi.org/10.7196/samj.2020.v111i1.14860>
12. Gholamrezanezhad A, Shirafkan A, Mirpour S, Rayatnavaz M, Alborzi A, Mogharrabi M, et al. Appropriateness of referrals for single-photon emission computed tomography myocardial perfusion imaging (SPECT-MPI) in a developing community: a comparison between 2005 and 2009 versions of ACCF/ASNC appropriateness criteria. *J Nucl Cardiol.* 2011;18(6):1044-52. Available from: <https://doi.org/10.1007/s12350-011-9419-3>
13. Gonzaga MA, Kiguli-Malwadde E, Businge F, Byanyima RK. Utilisation of obstetric sonography at a peri-urban health centre in Uganda. *Pan Afr Med J.* 2010;7(24):24. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3172624/>
14. Gorleku PN, Dzefi-Tetty K, Edzie EKM, Setorglo J, Piersson AD, Ofori IN, et al. The degree and appropriateness of computed tomography utilization for diagnosis of headaches in Ghana. *Heliyon.* 2021;7(4):e06722. Available from: <https://doi.org/10.1016/j.heliyon.2021.e06722>
15. Iñurrategui MC, Tablado MR, Esteban S, Kopitowski K, Marchitelli C, Terrasa S. Transvaginal ultrasound overuse in a private university hospital in Argentina: a cross-sectional study. *BMJ Evid Based Med.* 2022;27:A7-A8. Available from: <https://doi.org/10.1136/bmjebm-2022-podabstracts.14>
16. Jahanmehr N, Bigdeli AS, Salari H, Mokarami H, KhodaKarim S, Damiri S. Analyzing inappropriate magnetic resonance imaging (MRI) prescriptions and resulting economic burden on patients suffering from back pain. *Int J Health Plann Manage.* 2019;34(4):

- e1437-e1447. Available from: <https://doi.org/10.1002/hpm.2806>
17. Jame SZB, Majdzadeh R, Akbari Sari A, Rashidian A, Arab M, Rahmani H. Indications and overuse of computed tomography in minor head trauma. *Iran Red Crescent Med J*. 2014;16(5):e13067. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4082513/>
 18. Jame SZB, Sari AA, Majdzadeh R, Rashidian A, Arab M, Rahmani H. The extent of inappropriate use of magnetic resonance imaging in low back pain and its contributory factors. *Int J Prev Med*. 2014;5(8):1029-36. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4258662/>
 19. Kamrani R, Fallahi MJ, Masoompour SM, Ghayumi SMA, Jalli R, Khederzadeh S, et al. Evaluation of the appropriate use of chest CT-Scans in the diagnosis of hospitalized patients in shiraz teaching hospitals, Southern Iran. *Cost Eff Resour Alloc*. 2022;20(44). Available from: <https://doi.org/10.1186/s12962-022-00381-0>
 20. Kavosi Z, Sadeghi A, Lotfi F, Salari H, Bayati M. The inappropriateness of brain MRI prescriptions: a study from Iran. *Cost Eff Resour Alloc*. 2021;19(14). Available from: <https://doi.org/10.1186/s12962-021-00268-6>
 21. Kawooya M, Kisembo H, Malumba R, Nsereko EK. Effectiveness of clinical imaging guidelines to reduce inappropriate head computed tomography imaging: a case of Uganda. *Egypt J Radiol Nucl Med*. 53(164). Available from: <https://doi.org/10.1186/s43055-022-00833-w>
 22. Kawooya MG, Pariyo G, Malwadde EK, Byanyima R, Kisembo H. Assessing the performance of imaging health systems in five selected hospitals in Uganda. *J Clin Imag Sci*. 2012;2:12. Available from: <https://doi.org/10.4103/2156-7514.94225>
 23. Khiabani MS, Mohammadi MS, Ghoreyshi SA, Rohani P, Alimadadi H, Sohoul MH. Acute pancreatitis in 60 Iranian children: do pediatricians follow the new guidelines in diagnosis and management of acute pancreatitis? *BMC Pediatr*. 2022;22(457). Available from: <https://doi.org/10.1186/s12887-022-03509->
 24. Maitra D, Chatterjee S, Debnath S, Mukhopadhyay DK, Chattopadhyay S. Assessment of appropriateness of doing CT scan for investigating headache in a tertiary care hospital in eastern India. *Indian J Public Health Res Dev*. 2022;13(4):65-9. Available from: <https://doi.org/10.37506/ijphrd.v14i4.18531>
 25. Meidani Z, Hamidian Y, Farzandipour M, Aliasgharzade A. CT utilization: a case study in Iran based on ACR appropriateness criteria. *Radiol Manage*. 2017;39(1):33-38. Available from: <https://pubmed.ncbi.nlm.nih.gov/30725547/>
 26. Mensah YB, Mensah K, Gbadamosi H, Mensah NA. Magnetic resonance imaging (MRI) utilization in a Ghanaian teaching hospital: trend and policy implications. *Ghana Med J*. 2020;54(1): 3-9. Available from: <https://doi.org/10.4314/gmj.v54i1.2>
 27. Mohammadi N, Farahmand F, Hadizadeh Kharazi H, Mojdehipanah H, Karampour H, Nojomi M. Appropriateness of physicians' lumbosacral MRI requests in private and public centers in Tehran, Iran. *Med J Islam Repub Iran*. 2016;30:415. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5307604/#:~:text=In%20the%20private%20center%2C%2080.6,studie%20mostly%20from%20developed%20countries.>
 28. Pierson AD, Nunoo G, Gorleku PN. An audit of clinical practice, referral patterns, and appropriateness of clinical indications for brain MRI examinations: a single-centre study in Ghana. *Radiography*. 2018;24(2):e25-e30. Available from: <https://doi.org/10.1016/j.radi.2017.10.004>
 29. Refahi S, Kachooei AR, Farsadpour M, Shahrayeni R, Goudarzian M, Molavi Taleghani YM, et al. Is prescription of knee MRI according to standard clinical guideline? *Acta Med Mediterr*. 2016;4:1207-11. Available from: <https://www.actamedicamediterranea.com/archive/2016/special-issue-4/is-prescription-of-knee-mri-according-to-standard-clinical-guideline>
 30. Saadat S, Ghodsi SM, Firouznia K, Etminan M, Goudarzi K, Naieni KH. Overuse or underuse of MRI scanners in private radiology centers in Tehran. *Int J Technol Assess Health Care*. 2008;24(3):277-81. Available from: <https://doi.org/10.1017/S0266462308080379>
 31. Salari H, Omranikhoo H, Amini A, Amiri M, Bayyenas S, Azmal M, et al. Examining the amount of unnecessary knee MRI prescription in the MRI Center of Bushehr University of Medical Sciences in 2018. *Evid Based Health Policy, Manag Econ*. 2020;4(2):82-88. Available from: <https://doi.org/10.18502/jebhpme.v4i2.3433>
 32. El Sibai K, Itani S, Rabah A, Hourani M, Dakik HA. Evaluation of the appropriateness criteria for coronary computed tomography angiography in an academic medical center in a developing country: comparison of the 2006 and 2010 criteria. *J Nucl Cardiol*. 2011;18(6):1053-1058. Available from: <https://doi.org/10.1007/s12350-011-9437-1>
 33. Silva HS, Oliveira FKF, Prado LOM, Almeida-Santos M, Reis FP. Abdominal computed tomography in the emergency room: overuse of medical technologies

- and the depreciation of clinical diagnosis. *Rev Bras Educ Med.* 2019;43(Suppl 1). Available from: <https://doi.org/10.1590/1981-5271v43suplemento1-20190022>.ing
34. Sodhi KS, Krishna S, Saxena AK, Sinha A, Khandelwal N, Lee EY. Clinical application of "Justification" and "Optimization" principle of ALARA in pediatric CT imaging: "How many children can be protected from unnecessary radiation?" *Eur J Radiol.* 2015;84(9):1752-7. Available from: <https://doi.org/10.1016/j.ejrad.2015.05.030>
 35. Srivastava MK, Pagala RM, Kendarla V, Nallapareddy K. Appropriate use criteria in myocardial perfusion imaging in a tertiary care hospital in South India: an audit. *World J Nucl Med.* 2021;20(03):281-5. Available from: https://doi.org/10.4103/wjnm.wjnm_77_20
 36. Tambe J, Mbuagbaw L, Nguetack-Tsague G, Foyet J, Ongolo-Zogo P.. Multidetector computed tomography utilization in an urban sub-Saharan Africa setting: user characteristics, indications and appropriateness. *Pan Afr Med J.* 2020;37(42). Available from: <https://doi.org/10.11604/pamj.2020.37.42.21176>
 37. Tambe J, Onana Y, Dongmo S, Nguetack-Tsague G, Ongolo-Zogo P. Health insurance ownership and quality of computed tomography requests: experience from a peripheral referral hospital in Cameroon. *Radiol Res Pract.* 2021:e9959114. Available from: <https://doi.org/10.1155/2021/9959114>
 38. Tramuja L, Judice MM, Becker AB. Evaluation of the diagnostic management of deep vein thrombosis in the emergency department of a tertiary hospital in Santa Catarina, Brazil: a cross-sectional study. *J Vasc Bras.* 2022;21. Available from: <https://doi.org/10.1590/1677-5449.202002171>
 39. Veličković J, Hajdarević S, Palibrk IG, Janić NR, Đukanović M, Miljkovic B, et al. Routine chest radiographs in the surgical intensive care unit: can we change clinical habits with no proven benefit? *Acta Chir Iugosl.* 2013;60(3):39-44. Available from: <https://doi.org/10.2298/aci1303039v>
 40. Yu L, Wang X, Lin X, Wang Y. The use of lumbar spine magnetic resonance imaging in Eastern China: appropriateness and related factors. *PLoS One.* 2016;11:e0146369. Available from: <https://doi.org/10.1371/journal.pone.0146369>